

UNCLASSIFIED

AD NUMBER
AD854130
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; MAY 1969. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Releases Branch/TID, Frederick, MD 21701.
AUTHORITY
Army Biological Defense Research Lab ltr dtd 29 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD

TECHNICAL MANUSCRIPT 533

AD854130

NITROSOGUANIDINE MUTAGENESIS
AND CHROMOSOMAL REPLICATION
IN STAPHYLOCOCCUS AUREUS

Robert A. Altenbern

MAY 1969

JUN 20 1969

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

WHITE SECTION	<input type="checkbox"/>
BUFF SECTION	<input checked="" type="checkbox"/>
ANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
DISTRIBUTION/AVAILABILITY CODES	
DIST.	AVAIL. AND/OR SPECIAL
2	

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 533

NITROSOGUANIDINE MUTAGENESIS AND CHROMOSOMAL REPLICATION
IN STAPHYLOCOCCUS AUREUS

Robert A. Altenbern

STATEMENT #2 UNCLASSIFIED

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Dept. of Army, Fort Detrick, ATTN: Technical Release Branch/TID. Frederick. Maryland 21701

Medical Bacteriology Division
BIOLOGICAL SCIENCES LABORATORIES

Project 1B061102B71A

May 1969

NITROSOGUANIDINE MUTAGENESIS AND CHROMOSOMAL REPLICATION
IN STAPHYLOCOCCUS AUREUS

ABSTRACT

The number of mutations induced by nitrosoguanidine per 10^6 survivors is the same for exponential cells as for cells with resting, complete chromosomes. During synchronous chromosomal replication, no peak of mutations per million survivors appears at the time of duplication of the genes examined. It was concluded that for Staphylococcus aureus the replicating region of the chromosome is no more susceptible to nitrosoguanidine mutation than is the resting chromosome.

I. INTRODUCTION*

Recent investigations in several laboratories have led to varied methods for determining gene locations on the chromosome of several organisms. Wolf, Newman, and Glaser¹ developed an elegant method for mapping the Escherichia coli K-12 chromosome; a considerably different method was devised by Helmstetter² for E. coli B/r. Cerda-Olmeda, Hanawalt, and Guerola³ were able to derive a map of E. coli by nitrosoguanidine mutation during synchronous chromosomal replication; our own efforts^{4,5} in mapping the Staphylococcus aureus chromosome involved similar methodology.

It was shown that the replication point in the E. coli chromosome was significantly more amenable to nitrosoguanidine mutagenesis than the non-replicating regions.³ The report of this phenomenon led me to search for parallel behavior in S. aureus. To date, the evidence indicates that the replicating point of the S. aureus chromosome is no more mutable by nitrosoguanidine than is the non-replicating portion of the chromosome. In this context, S. aureus appears to be qualitatively different from E. coli.

* This report should not be used as a literature citation in material to be published in the open literature. Readers interested in referencing the information contained herein should contact the author to ascertain when and where it may appear in citable form.

II. MATERIALS AND METHODS

Single auxotrophs of two strains (Smith and Maybush) of S. aureus were isolated following exposure of the parent type to nitrosoguanidine and were employed throughout this study. The synchronous chromosomal replication method for mapping the S. aureus genome was carried out as previously described⁵ except that Albimi Brucella broth* replaced trypticase soy broth. Total viable counts were determined on plates of trypticase soy agar. Mutagen-induced prototrophs were detected and counted on minimal medium whose composition has already been described.⁴

Resting cells with completed chromosomes were prepared as follows: 10 ml of overnight culture in Albimi Brucella broth were added to 100 ml of fresh medium of the same composition. The culture was shaken at 37 C for 3 hours, at which time phenyl ethyl alcohol was added (0.40%) and incubation was continued at 30 C for 2 hours without shaking in a water bath. At the end of this incubation, the cells had ceased DNA synthesis and presumably possessed completed chromosomes. A 5-ml sample of these cells was centrifuged, the cells were resuspended in 5 ml of saline containing 200 µg nitrosoguanidine per ml and incubated at 30 C for 20 minutes. After this mutagen exposure, the cells were diluted in saline and plated on minimal agar and on trypticase soy agar to determine the number of induced reversions to prototrophy and the total survivors, respectively. Exponential cells were grown for 5 hours on a shaker at 37 C following inoculation of 5 ml of Albimi Brucella broth with 0.05 ml of an overnight culture in the same medium. These cells were then centrifuged and subjected to the same mutagen exposure as the resting cells.

III. RESULTS

The results of determination of the mutations per 10^6 survivors for three loci in each of two strains for exponential cells and for cells with resting chromosomes are presented in Table 1. It is clear that in strain Maybush the number of mutants per 10^6 survivors for exponential cells is the same as that for resting cells at each of the three loci. In strain Smith the mutations per 10^6 survivors in resting cells seem to be consistently higher than for exponential cells. These data contrast markedly with those reported for E. coli,³ in which the mutants per 10^6 survivors for exponential cells were 4 to 5 times greater than the same parameter determined for cells with resting chromosomes.

* Albimi Laboratories, Flushing, New York.

TABLE 1. FREQUENCY OF MUTATIONS PER 10^6 SURVIVORS INDUCED IN EXPONENTIAL CELLS AND CELLS WITH RESTING CHROMOSOMES FOR TWO STRAINS OF S. AUREUS

	<u>Mutations per 10^6 Survivors</u> ^{a/}		
	pan	rib	trp
<u>Strain Maybush</u>			
Exponential	13,100	6.8	10.9
Resting	14,680	5.3	10.1
<u>Strain Smith</u>			
Exponential	36	0.75	156
Resting	55	1.9	210

a. pan = pantothenate; rib = riboflavin;
trp = tryptophan.

Determination of total mutants, total survivors and, thus, mutants per 10^6 survivors for each of three loci for strain Smith during synchronous chromosomal replication led to the results presented in Figure 1. There is no peak of mutants per 10^6 survivors at the time of duplication of the gene in question. Total survivors seem to increase slightly with time, although other earlier data⁵ showed that there is no increase in number of cells before exposure to nitrosoguanidine.

The point in time of pronounced increase in mutants per 10^6 survivors at various loci in E. coli during synchronous chromosomal replication,³ designated the gene duplication time, has not been observed in S. aureus. The reasons for such a sharp contrast between these two organisms can be at present only a matter of conjecture. Cerda-Olmeda et al.³ speculated that, in E. coli, opening of the double-stranded chromosome during replication increased the mutability of that open region. In S. aureus this region does not exhibit increased mutability and may be partially protected by the replicase itself or by other aspects of the mechanics of replication.




FIGURE 1. Total Mutants and Total Survivors During Synchronous Chromosomal Replication Mapping of Auxotrophs of *S. aureus*, strain Smith.

LITERATURE CITED

1. Wolf, B.; Newman, A.; Glaser, D.A. 1968. On the origin and direction of replication of the Escherichia coli K12 chromosome. J. Mol. Biol. 32:611-629.
2. Helmstetter, C.E. 1968. Origin and sequence of chromosome replication in Escherichia coli B/r. J. Bacteriol. 95:1634-1641.
3. Cerda-Olmeda, E.; Hanawalt, P.C.; Guerola, N. 1968. Mutagenesis of the replication point by nitrosoguanidine: Map and pattern of replication of the Escherichia coli chromosome. J. Mol. Biol. 33: 705-719.
4. Altenbern, R.A. 1966. Apparent genomic mapping of Staphylococcus aureus by a new method. Biochem. Biophys. Res. Commun. 25:346-353.
5. Altenbern, R.A. 1968. Chromosome mapping in Staphylococcus aureus. J. Bacteriol. 95:1642-1646.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION
Department of the Army Fort Detrick Frederick, Maryland 21701		Unclassified
3. REPORT TITLE		2b. GROUP
NITROSOGUANIDINE MUTAGENESIS AND CHROMOSOMAL REPLICATION IN <u>STAPHYLOCOCCUS AUREUS</u>		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (First name, middle initial, last name)		
Robert A. Altenbern		
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
May 1969	11	5
8a. CONTRACT OR GRANT NO.	8b. ORIGINATOR'S REPORT NUMBER(S)	
a. PROJECT NO. 1B061102B71A	TECHNICAL MANUSCRIPT 533	
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		
10. DISTRIBUTION STATEMENT		
Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY	
	Department of the Army Fort Detrick Frederick, Maryland 21701	
13. ABSTRACT		
<p>The number of mutations induced by nitrosoguanidine per 10^6 survivors is the same for exponential cells as for cells with resting, complete chromosomes. During synchronous chromosomal replication, no peak of mutations per million survivors appears at the time of duplication of the genes examined. It was concluded that for <u>Staphylococcus aureus</u> the replicating region of the chromosome is no more susceptible to nitrosoguanidine mutation than is the resting chromosome.</p>		
14. Key Words		
<p>*<u>Staphylococcus aureus</u> *Nitrosoguanidine *Chromosomes Replicating Mutations <u>Escherichia coli</u> Auxotrophs</p>		

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

Unclassified

Security Classification